

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce; but at least one sample	Multi-methods													Additional analysis	
			Dithiocarbamates	Inorganic total bromide	Nitrate	Chlormequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphoric acid	Sulphur dioxide (SO ₂)		Heavy metals (Cadmium, Lead)
1. FRUITS FRESH OR FROZEN; NUTS																	
i) Citrus fruit																	
Grapefruit (Shaddocks, pomelos, sweeties, tangelo, ugli and other hybrids)	3	875	x							O(2, 4-D)	x	O	O	O			Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Grapefruit (Shaddocks, pomelos, sweeties, tangelo, ugli and other hybrids) (China, South Africa, Turkey)	7	170	x							O(2, 4-D)	x	O	O	O			Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Oranges	2	1600	x					x*		O(2, 4-D)	x	O	O	O			*Obligatory: third countries Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Lemons	2	1600	x					O*		O(2, 4-D)	x	O	O	O			Recommendation: for every 10th sample. *Recommendation: Overseas goods: ethephon at the beginning of the season Obligation: Sampling of each 4th sample after all process steps ²
Lemons (South America)	4	640	x					O*		O(2, 4-D)	x	O	O	O			
Lemons (South Africa)	5	480	x					O*		O(2, 4-D)	x	O	O	O			
Lemons (Turkey)	6	340	x					O*		O(2, 4-D)	x	O	O	O			
Limes	7	170	x							O(2, 4-D)	x	O	O	O			Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Mandarins (clementine, tangerine and other hybrids)	2	1600	x							O(2, 4-D)	x	O	O	O			Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
Mandarins (clementine, tangerine and other hybrids) (Third countries)	5	480	x							O(2, 4-D)	x	O	O	O			
Other citrus fruits	7	170	x							O(2, 4-D)	x	O	O	O			Recommendation: for every 10th sample Obligation: Sampling of each 4th sample after all process steps ²
ii) Tree nuts (shelled or unshelled)																	
Almonds	1	2500	x								x						
Brazil nut	1	2500	x								x						
Cashew nut	1	2500	x								x						
Chestnut	3	875	x	x*							x						*Obligation: Chloride/bromide-ratio
Chestnut (China, Turkey)	6	340	x	x*							x						Obligation: Sampling after all process steps
Coconut	1	2500	x								x						
Hazelnut	3	875	x								x						
Macadamia nut	1	2500	x								x						
Pecans	1	2500	x								x						

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ... t QS purchased produce; but at least one sample	Multi-methods	Dithiocarbamates	Inorganic total bromide	Nitrate	Chlormequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead)	Maleic hydrazide	Glyphosate	Diquat / Paraquat	Additional analysis	
Cultivated Blueberry	3	875	x								x											
Cowberry	5	480	x								x											
Cranberry	2	1600	x								x				x							
Currant (red, black and white)	6	340	x								x											
Gooseberry	7	170	x								x											
Jostaberry	3	875	x								x											
Rose hip	3	875	x								x				x							
Mulberry	3	875	x								x				x							
Elderberries (wild rowan berry)	3	875	x								x											
Cape gooseberry; Physalis	2	1600	x								x											
Other small fruits and berries	7	170	x								x				x							
vi) Miscellaneous fruit																						
a) Edible peel																						
Date	1	2500	x								x				x*							*Obligation: for each 3rd sample
Fig	1	2500	x						x**		x				x*							*Obligation: for each 3rd sample
Fig (Brazil)	4	640	x						x**		x				x*							**Obligation: for origin Brazil, for origin Turkey at the beginning of the season until end of August
Table olive	5	480	x								x				x*							*Obligation: for each 3rd sample
Kumquat	6	340	x						x**		x				x*							*Obligation: for each 3rd sample
Carambola	7	170	x								x				x*							**Obligation: at the beginning of the season
Kaki; Japanese persimmons	3	875	x						x*		x											*Obligation: for each 3rd sample
Other miscellaneous fruits with edible peel	7	170	x								x											*Obligation: for origin Spain at the beginning of the season
b) Inedible peel, small																						
Kiwi	1	2500	x								x											
Kiwi (Chile)	4	640	x								x											
Lychee	4	640	x								x											
Maracuja; Passionfruit (Granadilla)	9	45	x								x				x	x						
Prickly pear; cactus fruit; pitaya	8	90	x								x				x							
Other small miscellaneous fruits with inedible peel	9	45	x								x				x							
c) Inedible peel, large																						
Avocado	4	640	x								x											
Avocado (Chile, Peru)	6	340	x								x					x*						*Obligation: for each 4th sample
Banana	2	1600	x								x				x							
Mango	5	480	x								x				x							
Mango (Brazil)	7	170	x								x				x							
Papaya; Tamarillo	7	170	x								x				x							
Pomegranate	5	480	x								x				x							
Pomegranate (Turkey)	9	45	x								x				x							
Pineapple	1	2500	x						x		x											
Bread fruit; Jackfruit	6	340	x								x				x							
Cherimoya; Rambutan	6	340	x								x				x							
Durian	6	340	x								x				x							
Guava	8	90	x								x				x							
Other large miscellaneous fruits with inedible peel	9	45	x								x				x							

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce; but at least one sample	Multi-methods														Additional analysis	
			Dithiocarbamates	Inorganic total bromide	Nitrate	Chlormequat / Mepiquat	Dithianon	Etthephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead)		Maleic hydrazide
2. VEGETABLES FRESH OR FROZEN																		
i) Root and tuber vegetables																		
a) Potatoes																		
Potato	1	2500	x	O														
b) Tropical root and tuber vegetables																		
Ginger	9	45	x															*Obligation: for each 3rd sample
Cassava (Dasheen, eddoe (Japanese taro), tannia, manioc)	4	640	x															*Obligation: for each 3rd sample
Sweet potato	1	2500	x															*Obligation: for each 3rd sample
Yams (Potato bean (yam bean), Mexican yam bean)	4	640	x															*Obligation: for each 3rd sample
Other tropical root and tuber vegetables	9	45	x															
(c) Other root and tuber vegetables except sugar beet																		
Beetroot	1	2500	x															*Obligation: for each 3rd sample
Carrot (bunch)	2	1600	x															*Obligation: for each 3rd sample
Carrot (bulk goods)	2	1600	x															
Carrot (bulk goods) (Germany, Netherlands, Spain)	1	2500	x															*Obligation: for each 3rd sample
Carrot (bulk goods) (Israel)	3	875	x															
Celeriac	6	340	x															*Obligation: for each 3rd sample
Horseradish	1	2500	x															*Obligation: for each 3rd sample
Jerusalem artichoke	1	2500	x															*Obligation: for each 3rd sample
Parsnip	3	875	x															*Obligation: for each 3rd sample
Parsley root	3	875	x															*Obligation: for each 3rd sample
Radish (Outdoor)	2	1600	x															
Radish (Greenhouse)	1	2500	x															
Small radish (Outdoor)	3	875	x															
Small Radish (Greenhouse)	1	2500	x															
Scorzonera	1	2500	x															
White Turnip; Turnip; Swedes	3	875	x															*Obligation: for each 3rd sample
Other root and tuber vegetables except sugar beet	6	340	x															
ii) Bulb vegetables																		
Garlic	1	2500	x															
Garlic (fresh)	4	640	x															
Onions (Silver onions)	1	2500	x															*Obligation: for each 3rd sample
Onions	1	2500	x															
Onions (Egypt, New Zealand)	2	1600	x															
Shallots	1	2500	x															*Obligation: for each 3rd sample
Spring onion	2	1600	x															
Spring onions (Egypt; Italy)	6	340	x															
Other bulb vegetables	6	340	x															
iii) Fruiting vegetables																		
a) Solanaceae																		
Tomato	1	2500	x															
Tomato (Spain, Italy)	2	1600	x															



Qualitätssicherung. Vom Landwirt bis zur Ladentheke.



Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per unit QS purchased produce; but at least one sample	Multi-methods	Dithiocarbamates	Inorganic total bromide	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphoric acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead)	Maleic hydrazide	Glyphosate	Diquat / Paraquat	Additional analysis
Tomato (Turkey)	3	875	x								x										

Product	Risk group ¹	Wholesale/Preparation and Processing: one sample per lot QS purchased produce; but at least one sample	Multi-methods	Dithiocarbamates	Inorganic total bromide	Nitrate	Chloromequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) ³	Fenbutatin oxide	QAV	Morpholine	Phosphoric acid	Sulphur dioxide (SO ₂)	Heavy metals (Cadmium, Lead)	Maleic hydrazide	Glyphosate	Diquat / Paraquat	Additional analysis
			x	x*																	
Tarragon (Hyssop)	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Wild garlic	8	90	x								x										
Caraway	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample

