





Product	Risk group <sup>1</sup>	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce; but at least one sample	Residue Monitoring											Additional analysis					
			Multi-methods Dithiocarbamates	Inorganic total bromide	Nitrate	Chlormequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) <sup>3</sup>	Fenbutatin oxide	QAV	Morpholine		Phosphonic acid Sulphur dioxide (SO <sub>2</sub> )	Heavy metals (Cadmium, Lead)	Maleic hydrazide	Glyphosate	Diquat / Paraquat
Currant (red, black and white)	6	340	x																
Gooseberry	7	170	x																
Jostaberry	3	875	x																
Rose hip	3	875	x																
Mulberry	3	875	x																
Elderberries (wild rowan berry)	3	875	x																
Cape gooseberry; Physalis	2	1600	x																
Other small fruits and berries	7	170	x																
<b>vi) Miscellaneous fruit</b>																			
<b>a) Edible peel</b>																			
Date	1	2500	x																*Obligation: for each 3rd sample
Fig	1	2500	x																*Obligation: for each 3rd sample
Fig (Brazil, Israel)	4	640	x						x**										**Obligation: for origin Brazil, for origin Turkey at the beginning of the season until end of August
Table olive	5	480	x																*Obligation: for each 3rd sample
Kumquat	6	340	x						x**										*Obligation: for each 3rd sample **Obligation: at the beginning of the season
Carambola	6	340	x																*Obligation: for each 3rd sample
Kaki; Japanese persimmons	3	875	x						x*										*Obligation: for origin Spain at the beginning of the season
Other miscellaneous fruits with edible peel	6	340	x																
<b>b) Inedible peel, small</b>																			
Kiwi	1	2500	x																
Kiwi (Chile)	3	875	x																
Lychee	3	875	x																
Maracuja; Passionfruit (Granadilla)	9	45	x																
Prickly pear; cactus fruit; pitaya	8	90	x																
Other small miscellaneous fruits with inedible peel	9	45	x																
<b>c) Inedible peel, large</b>																			
Avocado	4	640	x																
Avocado (Chile, Peru)	6	340	x																*Obligation: for each 4th sample
Banana	2	1600	x																
Mango	5	480	x																
Mango (Brazil, Dominican republic, Peru)	6	340	x																
Papaya; Tamarillo	7	170	x																
Pomegranate (EU, South Africa)	5	480	x																
Pomegranate (Third Countries)	9	45	x																
Pineapple	1	2500	x																
Bread fruit; Jackfruit	6	340	x																
Cherimoya; Rambutan	6	340	x																
Durian	6	340	x																
Guava	8	90	x																
Other large miscellaneous fruits with inedible peel	9	45	x																
<b>2. VEGETABLES FRESH OR FROZEN</b>																			
<b>i) Root and tuber vegetables</b>																			
<b>a) Potatoes</b>																			







Product	Risk group <sup>1</sup>	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce; but at least one sample	Multi-methods	Dithiocarbamates	Inorganic total bromide	Nitrate	Chlormequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) <sup>3</sup>	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO <sub>2</sub> )	Heavy metals (Cadmium, Lead)	Maleic hydrazide	Glyphosate	Diquat / Paraquat	Additional analysis
Celery leaves; Sorrel	8	90	x								x										
Coriander leaves	8	90	x								x										
Lovage	8	90	x								x										
Parsley	8	90	x								x										
Sage	8	90	x								x										
Rosemary	8	90	x								x										
Thyme	8	90	x								x										
Basil	8	90	x								x										
Mint	8	90	x								x										
Tarragon (Hyssop)	8	90	x								x										
Wild garlic	8	90	x								x										
Caraway	8	90	x								x										
Bay leaves	8	90	x								x										
Marjoram	8	90	x								x										
Oregano	8	90	x								x										
Savory	8	90	x								x										
Common balm; Lemon balm	8	90	x								x										
Edible flowers	8	90	x								x										
Other fresh herbs and edible flowers	9	45	x								x										
<b>Cut herbs</b>																					
<sup>(2)</sup> for entire g) fresh herbs (Italy, Third Countries)	9	45	x	x*							x										*Obligation: from November to March for each 4th sample
Chervil	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Chives	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Dill leaves	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Celery leaves; Sorrel	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Coriander leaves	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Lovage	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Parsley	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Sage	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Rosemary	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Thyme	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Basil	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample

Product	Risk group <sup>1</sup>	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce; but at least one sample	Multi-methods	Dithiocarbamates	Inorganic total bromide	Nitrate	Chlormequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) <sup>3</sup>	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO <sub>2</sub> )	Heavy metals (Cadmium, Lead)	Maleic hydrazide	Glyphosate	Diquat / Paraquat	Additional analysis
Mint	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Tarragon (Hyssop)	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Wild garlic	8	90	x								x										
Caraway	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Bay leaves	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Marjoram	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Oregano	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Savory	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Common balm; Lemon balm	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Edible flowers	8	90	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
Other fresh herbs and edible flowers	9	45	x	x*							x										*Obligation: for origin Southern Europe from November to March for each 4th sample
<b>vi) Legume vegetables (fresh)</b>																					
Beans (with pods)	2	1600	x								x										
Beans (with pods) (Spain)	4	640	x								x										
Beans (with pods) (Third Countries)	6	340	x								x										
Beans (without pods)	2	1600	x								x										
Beans (without pods) (Third Countries)	4	640	x								x										*Obligation: for each 3rd sample
Peas (with pods) (Mangetout (sugar peas))	3	875	x								x										*Obligation: for each 3rd sample
Peas (with pods) (Third Countries)	7	170	x								x										*Obligation: for each 3rd sample
Peas (without pods)	2	1600	x								x										*Obligation: for each 3rd sample
Peas (without pods) (Third Countries)	4	640	x								x										*Obligation: for each 3rd sample
Other legume vegetables (fresh)	7	170	x								x										
<b>vii) Stem vegetables (fresh)</b>																					





Product	Risk group <sup>1</sup>	Wholesale/Preparation and Processing: one sample per ...t QS purchased produce; but at least one sample	Multi-methods	Dithiocarbamates	Inorganic total bromide	Nitrate	Chlormequat / Mepiquat	Dithianon	Ethephon	Phenoxyalkane carboxylic acids (alkaline hydrolysis)	Matrine (Single method only required for positive findings from the multimethod) <sup>3</sup>	Fenbutatin oxide	QAV	Morpholine	Phosphonic acid	Sulphur dioxide (SO <sub>2</sub> )	Heavy metals (Cadmium, Lead)	Maleic hydrazide	Glyphosate	Diquat / Paraquat	Additional analysis
<b>5. PSEUDOCEREALS</b>																					
Pseudocereals (Amaranth, Buckwheat, Chia seeds, Quinoa)	8	90	x								x									x	x

**Legend:**

x  
\*  
O

obligatory analysis  
analysis with separate explanation  
additional recommendation

Multi methods:  
GC  
LC-MS/MS

Group of selected pesticides identified by gas chromatography  
Group of selected pesticides identified by LC-MS/MS

2,4-D

Herbicide from the group of Phenoxyalkyl carbonic acid

<sup>1</sup>The risk grouping of the products is made by QS and is based on a scale from 1 (lowest risk) to 9 (highest risk).

<sup>2</sup>At least the first sample and afterwards every 4th sample needs to be taken after all process steps

<sup>3</sup>For positive detections of Matrine by multi-method(s) a precise quantification of the active substance by a single method is required, if this is not ensured by the multi-methods.

**Revision Information Version 01.01.2021rev01 (as of 01.07.2021)**

Criterion/ Requirement	Changes	Date of change
10.1 Control plan	Change: The tonnage per risk group has been revised.	01.07.2021